

## **Genetic Diversity Analysis of Eighteen Tea (*Camellia sinensis* L.) Clones of Bangladesh Through RAPD**

**Shefali Boonerjee, M. Nurul Islam, M. I. Hoque and R.H. Sarker**

*Department of Botany, University of Dhaka, Dhaka-1000, Bangladesh*

*Key words: Camellia sinensis, RAPD, Genetic diversity, Cluster analysis, Dendrogram*

### **Abstract**

Using 20 decamer random primers molecular characterization of 18 tea (*Camellia sinensis* L.) clones of Bangladesh was made. All the primers showed significant amplification in PCR analysis. A total of 755 bands was produced in all the 18 tea clones with an average of 37.75 RAPD bands per primer. Among all the bands 97.41% were polymorphic in nature. The molecular size of the amplified DNA fragments ranged from 250 to 5000 bp. Ten unique bands were amplified from the genome of the 18 tea clones. The values of pairwise genetic distance ranged from 24.0 to 59.0 indicating the presence of a wide range of genetic diversity. The highest genetic distance 59 was found between the clone BT16 and BT2, whereas the lowest (24.0) between BT18 and BT5. The dendrogram based on Nei's genetic distance was constructed using un-weighted Pair Group of Arithmetic Mean (UPGMA) segregating the 18 tea clones into two major clusters: BT9 and BT13 in cluster 1 and the remainder of 16 clones in cluster 2. Cluster 2 is further subdivided into many sub-clusters. Cluster analysis revealed that while the genotype BT5 is closely related to BT18, BT1 and BT2 showed similarity with BT8. Genotypes BT1 and BT13 were widely diverse genetically.

### **Introduction**

Tea (*Camellia sinensis* L.) is the oldest mild stimulating drink containing caffeine used worldwide. Tea leaves have more than 700 chemical constituents; among them flavanoides, amino acids, vitamins (B, C, E, K), caffeine and polysaccharides are important to human health (Mondal et al. 2004). It is being used as an important health drink from the ancient time and recent literature strongly emphasizes its value in disease prevention and as a key component of daily diets (Mukhtar and Ahmad 1999, 2000 and Weisburger 1997). Bangladesh is a tea-exporting country and the tea industry is one of the most important labor and export oriented industries in Bangladesh. Bangladesh earns a substantial amount

of foreign exchange contributing to 11% of the GDP (BTB 2012). At present the global scenario of tea market is very competitive and expected to be more in the near future. Specific cultivar-based made tea might be the requirement of the future generation of tea consumers, whereas tea industries in Bangladesh are now facing an international competition because of its low quality. Increase in per unit production as well as improvement in quality is important for the survival of tea industries in Bangladesh. Widespread natural hybridization increases genetic variability in tea which has been usually exploited for developing many new cultivars of Bangladesh. Exploiting the genetic variability present in the natural population, the Bangladesh Tea Research Institute (BTRI) has developed 18 recommended high yielding and qualitative clones since the last 55 years. These clones were classified and identified traditionally on the basis of morphological traits only on the leaf morphology. Due to phenotypic plasticity the germplasm may show different morphology Goodrich et al. (1985). So because of environmental influences true estimation of genetic diversity for characterization and cataloguing the clones using morphological distinctiveness is difficult. However, genetic information of the clones is the basic requirement for improvement using hybridization. Accurate and authentic data at the DNA level are important for the assessment of genetic diversity and to study the phylogenetic relationship of these distinctive clones. But no such information is available for these clones. At present DNA based molecular markers (RFLPs, SSRs and RAPDs) are effective and reliable for measuring genetic diversity and evolutionary relationship among crop germplasms (Kidwell et al. 1994 and Mengoni et al. 2000).

Among the DNA markers, development of RAPD-PCR based DNA finger printing is easier (Gherardi et al. 1998). Since the discovery of RAPD assay is being used in a number of areas of plant taxonomy (Williams et al. 1990). Through this technique several crop plant germplasms like coffee (Orozco-Castillo et al. 1994), rice (Virk et al. 1995), mungbean (Sony et al. 2012, Saini et al. 2010, Lakhanpaul et al. 2000), *Cicer arietinum* L. and *Cajanus cajan* L. (Datta and Lal 2011), tomato (Tabassum et al. 2013, Gubba and Sivparsad 2008, El-Hady et al. 2010 and Ezekiel et al. 2011) and many more have been characterized.

Molecular characterization and genetic documentation of tea germplasm including distinctive cultivars using RAPD have been reported earlier (Wachira et al. 1995, Bera and Saikia 2002, Mondal et al. 2004, Balasaravanan et al. 2003, Mishra and Mandi 2004, Mishra et al. 2009). But no such study for characterization and genetic diversity analysis of Bangladeshi tea clones was done before.

Therefore, the objective of the present study was to investigate and compare the RAPD fingerprinting-based genetic diversity among 18 tea clones released from the Bangladesh Tea Research Institute (BTRI).

## Materials and Methods

Leaf samples of 18 Bangladesh Tea (BT) clones, namely, BT1 - 18 were collected from the nucleus clone plot (NCP) of BTRI, Srimangal. Information about the clones used in this study is shown in Table 1.

**Table 1. Basic information regarding the 18 tea clones used for estimation of genetic diversity.**

Sl. No.	Name of the accessions	Name of the clones	Type
1	B201/39	BT1	Hybrid 2
2	R5/8	BT2	Hybrid 2
3	U16/21	BT3	Hybrid 3
4	B24/44	BT4	Manipuri Hybrid
5	BS1/2	BT5	Hybrid-3
6	BS1/1	BT6	Hybrid 3
7	Br/2/99	BT7	Manipuri
8	B90/44	BT8	Manipuri Hybrid
9	Balu 2/4	BT9	Hybrid 2
10	D/13/25	BT10	Assam
11	TV/E	BT11	Hybrid-3
12	H1/15	BT12	Manipuri
13	SH/D/11/333	BT13	Assam
14	B/HB/6/4	BT14	Manipuri-Hybrid
15	BS1/1/111	BT15	Hybrid-3
16	SH/D/11/313	BT16	Assam
17	T1B2	BT17	Hybrid 2
18	A/22/39	BT18	Assam

Total genomic DNA was extracted by using modified CTAB method (Doyle and Doyle 1987). DNA concentration was quantified through spectrophotometer (Simadzu Corp. Japan). The A260/280 readings for DNA samples were 1.6-1.8.

The PCR reaction mixture was prepared for 25 µl containing template DNA (25 ng) 2 µl, de-ionized distilled water 18.8 µl, Taq buffer A 10x (10 mM Tris-HCl with 1.5 mM MgCl<sub>2</sub>) 2.5 µl, primer (10 µM) 1.0 µl, dNTP mix (10 mM) 0.5 µl, Taq DNA polymerase (5 U/µl) 0.2 µl. PCR amplification was done in an oil-free thermal cycler (Biometra UNOII, Germany) for 46 cycles after initial denaturation at 94°C for 5 min and at 94°C for 1 min, annealing at 36°C for 30 sec. extension at 72°C for 3 min and final extension at 72°C for 5 min.

Twenty random decamer primers were used in the present study which showed reproducible results (Table 2). The amplified products were separated by horizontal electrophoresis using 1% agarose gel containing 0.5 µg/ml ethidium

bromide in TAE buffer at 50 volts and 100 mA for 1.5 h. A 1.0 kb ladder was electrophoresed alongside RAPD analysis as a marker. The gel was visualized by UV-transilluminator to examine the banding patterns and photographed by gel documentation system (CSL-Microdoc System, Cleaver Scientific Ltd. USA).

For analysis of genetic diversity the photographs of RAPD gel were critically examined on the basis of presence (1) or absence (0) of bands, size of bands and overall polymorphism of bands. The scores obtained using all primers in the RAPD analysis were then pooled for constructing a single data matrix. This was used for estimating polymorphic loci, genetic diversity, genetic distance (D) and constructing UPGMA (Sneath and Sokal 1973) dendrogram among the clones using computer program "Statistica".

## Results and Discussion

The present study was conducted to reveal the genetic variation among the 18 tea clones by reproducible amplification of DNA through RAPD analysis using 20 random decamer primers. The primer sequences used for amplification are enumerated in Table.2. A total of 755 distinct bands was scored from 20 different fingerprinting profiles of which 737 were polymorphic. An average of 37.75 bands per primer was amplified showing 97.41% polymorphic amplification which indicates the high level of polymorphism present among the 18 tea clones. A diverse level of polymorphism in different crops has been reported in *Brassica* (Chen et al. 2000), tomato (Moonmoon 2006, Tabassum et al. 2013), egg plant (Biswas et al. 2009) and chili (Paran et al. 1998). Extensive polymorphism in tea was reported earlier in several genetic diversity studies using RAPD markers (Wachira et al. 1995, Kaundun et al. 2000, Jorge et al. 2003). Mondal et al. (2000) characterized 25 Indian and 2 commercial tea cultivars using RAPD markers and found 95.2% genetic variability. The data set produced in this experiment was sufficient to categorize all the 18 tea clones (Table 2). Among the 20 primers OPA-7 produced maximum number of polymorphic bands (95 bands) that indicated a high level of polymorphism. On the other hand primer B06 generated the least number of polymorphic bands (14 bands). The RAPD banding profiles of the amplified products of three representative primers are shown in Figs. 1, 2 and 3. The band size ranged from 250 to 5000 bp. In earlier reports, Lai et al. (2001) and Gul et al. (2007) observed approximately 200 - 1000 bp and 250 - 1500 bp fragment sizes in different tea clones respectively.

Moreover, ten clone-specific bands were identified for different primer in different clones. These specific bands can successfully be used to generate genetic markers for identification of clones. Primer OPA 4, OPA 6, OPA 9, C01, C02, A03 and B14 were found to be the most effective to generate unique bands (Table 2).

**Table 2.** RAPD fingerprinting profile with 20 primers in 18 tea clones.

Primer code	Sequence (5'-3')	No. of amplified buds	Band size range (bp)	Unique bands with clones and bp size	No. of polymorphic bands
OPA-1	TGC CGA GCT C	59	500 - 2000	-	59
OPA-2	TGC CGA GCT G	52	250 - 2000	-	52
OPA-3	AGT CAG CCA C	38	500 - 1500	-	38
OPA-4	AAT CGG GCT G	15	750 - 3500	BT2 (3500)	15
OPA-5	AGG GGT CTT G	24	1500 - 3500	-	24
OPA-6	GGT CCC TGA C	42	250 - 3000	BT17 (250)	42
OPA-7	GAA ACG GGT G	95	250 - 3000	-	95
OPA-8	GTG ACG TAG G	39	250 - 1500	-	21
OPA-9	GTG ATC GCA G	27	750 - 3000	BT10 (750)	27
OPA-10	GTG ATC GCA G	28	750 - 2000	-	28
A 15	TTC CGA ACC C	45	1000 - 4000	-	45
A09	GGG TAA CGC C	27	750 - 5000	BT3 (5000) BT7 (750)	27
C01	TTC GAG CCA G	46	750 - 4000	BT2 (750)	46
C02	GTG AGG CGT C	71	750 - 4500	BT3 (4500)	71
D02	GGA CCC AAC C	19	1000 - 3500	-	19
A03	AGT CAG CCA C	33	250 - 3500	BT7 (3500) BT7 (250)	33
A08	AGT CAG CCA C	42	500 - 5000	-	42
D01	ACC GCG AAG G	24	750 - 4000	-	24
B14	TCC GCT CTG G	15	1000 - 3500	BT16 (3000)	15
B06	TGC TCT GCC C	14	2000 - 4000	-	14
Total		755			737




Fig. 1. RAPD profiles with primer OPA-6 in 18 clones of tea. M : 1 kb DNA ladder, 1: BT1, 2: BT2, 3: BT3, 4: BT4, 5: BT5, 6: BT6, 7: BT7, 8: BT8, 9: BT9, 10: BT10, 11: BT11, 12: BT12, 13: BT13, 14: BT14, 15: BT15, 16: BT16, 17: BT17 and 18: BT18.

The values of pairwise Nei's (1972) genetic distance ranged from 24.00 to 59.00. According to the distance matrix the highest genetic distance (59.00) were found between BT17 and BT3 while that of lowest (24) was in BT18 and BT5 (Table 3). The difference between the highest and lowest value of genetic distance revealed a wide range of variability persisting among the 18 selected tea

clones. From the result of genetic dissimilarity analysis using UPGMA method Gul et al. (2007) reported an extensive genetic diversity in the 24 tea genotypes. Welsh and MacClelland (1990) and Dos Santos et al. (1994) also confirmed the wide range of genetic diversity among the tea genotypes from their study through RAPD analysis.




Fig. 2. RAPD profiles with primer OPA-7 in the 18 clones of tea. M : 1 kb DNA ladder on the extreme left, 1: BT1, 2: BT2, 3: BT3, 4: BT4, 5: BT5, 6: BT6, 7: BT7, 8: BT8, 9: BT9, 10: BT10, 11: BT11, 12: BT12, 13: BT13, 14: BT14, 15: BT15, 16: BT16, 17: BT17 and 18: BT18.




Fig 3. RAPD profiles with primer C02 in the 18 clones of tea. M : 1 kb DNA ladder, on the extreme left 1: BT1, 2: BT2, 3: BT3, 4: BT4, 5: BT5, 6: BT6, 7: BT7, 8: BT8, 9: BT9, 10: BT10, 11: BT11, 12: BT12, 13: BT13, 14: BT14, 15: BT15, 16: BT16, 17: BT17 and 18: BT18.




Fig. 4. UPGMA dendrogram based on Nei's genetic distance summarizing the data on differentiation among 18 tea clones as revealed by RAPD analysis.

Table 3. Summary of Nei's genetic distances in 18 tea clones.

Variable	BT1	BT2	BT3	BT4	BT5	BT6	BT7	BT8	BT9	BT10	BT11	BT12	BT13	BT14	BT15	BT16	BT17	BT18	
BT1	*																		
BT2	32.0	*																	
BT3	36.0	46.0	*																
BT4	38.0	42.0	48.0	*															
BT5	39.0	31.0	45.0	37.0	*														
BT6	45.0	47.0	39.0	45.0	36.0	*													
BT7	37.0	53.0	43.0	43.0	48.0	52.0	*												
BT8	38.0	42.0	46.0	46.0	43.0	45.0	51.0	*											
BT9	46.0	36.0	48.0	42.0	41.0	53.0	53.0	52.0	*										
BT10	51.0	49.0	57.0	47.0	34.0	34.0	54.0	47.0	49.0	*									
BT11	45.0	51.0	45.0	45.0	40.0	44.0	46.0	45.0	53.0	48.0	*								
BT12	44.0	38.0	54.0	42.0	31.0	51.0	53.0	44.0	44.0	51.0	39.0	*							
BT13	39.0	43.0	57.0	45.0	46.0	52.0	42.0	49.0	41.0	52.0	46.0	49.0	*						
BT14	39.0	51.0	43.0	37.0	40.0	52.0	42.0	55.0	41.0	50.0	44.0	43.0	42.0	*					
BT15	49.0	49.0	49.0	43.0	44.0	48.0	52.0	51.0	51.0	46.0	38.0	41.0	48.0	46.0	*				
BT16	49.0	59.0	49.0	53.0	44.0	46.0	52.0	53.0	59.0	44.0	38.0	47.0	48.0	38.0	42.0	*			
BT17	43.0	41.0	59.0	41.0	28.0	48.0	48.0	47.0	39.0	40.0	42.0	35.0	46.0	44.0	30.0	44.0	*		
BT18	37.0	41.0	45.0	41.0	24.0	44.0	32.0	45.0	51.0	42.0	36.0	37.0	42.0	30.0	38.0	34.0	28.0	*	

The dendrogram constructed based on Nei's (1972) genetic distance separated out the 18 clones into two major clusters (Fig. 2). Cluster 1 contained 2 clones and the remainder 16 in cluster 2. BT9 and BT13 formed cluster 1 showing the close relatedness between them. Cluster 2 was further sub-divided into 2 sub-clusters forming more sub-sub clusters. BT1, BT2, BT8, BT3 and BT7 were separately grouped in sub-cluster 1. On the other hand, sub-cluster 2 comprises four separate groups. BT6 and BT10 made group 1 in this sub-cluster, while BT16 alone made an individual group 2. In group 3, clone BT15, BT11, BT12, BT17, BT18 and BT5 were included, whereas BT4 and BT14 in group 4. On the basis of cluster analysis it was observed that clone BT5 was closely related to BT18 according to the phylogenetic relationship as they maintained lowest genetic distances (24.00) among the 18 members. BT1 and BT2 were closely related showing similarity with BT8. Close relationship was observed among the pair of BT18 and BT17, BT12 and BT17, BT6 and BT10. Placing in the same cluster BT5 and BT10 maintained distant relationship. Though the clones were grouped in the same cluster, they were distantly related with each other.

From this investigation it is clearly indicating that each of the 18 tea clones possessed specific fingerprinting profile which could be used for their authentic identification. The detection of high level of genetic variability also supports the findings of Lai et al. (2001), Chen et al. (2000), and further strengthens the reports of Welsh and MacClelland (1990) and Dos Santos et al. (1994) who concluded that RAPD markers are effective for visualizing high level of genetic polymorphism in plant species.

### Acknowledgements

The study is a part of the NATP phase-1 Ph.D. research program funded by Bangladesh Agricultural Research Council (BARC). The authors are grateful to Professor M. Mozammel Haque of the Department of Botany, University of Dhaka for his kind help during the preparation of this manuscript.

### References

- Balasaravanan T, Pius PK, Rajkumar R, Muraleedharan N and Shasany AK** (2003) Genetic diversity among South Indian tea germplasm (*Camellia sinensis*, *C. assamica* and *C. assamica* spp. *lasicalyx*) using ALFP markers. *Plant Science* **165**: 365-375.
- Bangladesh Tea Board** (2012) Statistics on tea. Bangladesh Tea Board, Nasirabad, Chittagong.
- Bera B and Saikia H** (2002) Randomly amplified polymorphic DNA (RAPD) marker analysis in tea (*Camellia sinensis* L.) generative clones. *In: Proc. Plantation Crops Symposium-XV on "Sustainability of Plantation Crops through Integrated Approaches for Crop Production and Product Diversification"* held at the Central Coffee Research Institute during 10-13 December, 2002, Mysore, Karnataka, pp. 235-238.



- Biswas MS, Akhond AAY, Alamin M, Khatun M and Kabir MR** (2009) Genetic relationship among ten promising eggplant varieties using RAPD markers. *Plant Tissue Cult. & Biotech.* **19**(2): 119-126.
- Chen YP, Cao-Jia S, Miao Y, Cao JS and Ye WZ** (2000) Analysis of genetic polymorphisms in vegetable crops of *Brassica campestris* by RAPD markers. *J. Zhejiang University. Agric. Life Sci.* **26**: 131-136.
- Datta J and Lal N** (2011) Characterization of genetic diversity in *Cicer arietinum* L. and *Cajanus cajan* L. Millspaugh using random amplified polymorphic DNA and simple sequence repeat markers. *Genom. and Quant. Genet.* **3**: 30-41.
- Dos Santos JB, Nienhuis J, Skroch P, Tivang J and Slocam** (1994) Comparison of RAPD and RELP genetic markers in determining genetic similarity in *Brassica oleracea* L. genotypes. *TAG.* **87**: 909-915.
- Doyle JJ and Doyle JL** (1987) A rapid DNA isolation procedure from small quantities of fresh leaf tissues. *Phytochem. Bull.* **96**: 11-15.
- El-Hady EAAA, Haiba AAA, EL-Hamid NRA and Rizkalla AA** (2010) Phylogenetic diversity and relationships of some tomato varieties by electrophoretic protein and RAPD analysis. *J. Amer. Sci.* **6**(11): 434-441.
- Ezekiel CN, Nwangburuka CC, Ajibade OA and Odebode AC** (2011) Genetic diversity in 14 tomato (*Lycopersicon esculentum* Mill.) varieties in Nigerian markets by RAPD-PCR technique. *Afr. J. Biotechnol.* **10**(25): 4961-4967.
- Gherardi M, Mangin B, Goffinate B, Bonnet D and Huguet T** (1998) A method to measure genetic distance between allogamous population of alfalfa (*Medicago sativa*) using RAPD molecular markers. *TAG* **96**: 406-412.
- Goodrich WJ, Cook RJ and Morgan AG** (1985) The application of electrophoresis to the characterization of cultivars *Vicia faba* L. *Fabis Newsletter* **13**:8.
- Gubba A and Sivparsad BJ** (2008) Isolation and molecular characterization of tomato spotted wilt virus (TSWV) isolates occurring in South Africa. *Afr. J. Agri. Res.* **3**(6): 428 - 434.
- Gul S, Ahmed H, Khan IA and Alam M** (2007) Assessment of genetic diversity in tea genotypes through RAPD primers. *Pakistan J. Biol. Science.* **10**(15): 2609-2611.
- Jorge S and Pedroso MC** (2003) Genetic Differentiation of Portuguese Tea Plant using RAPD Markers. *Hort Science* **38**(6): 1191-1197.
- Kaundan SS, Zhyvoloup A and Park Y** (2000) Evaluation of the genetic diversity among elite tea accessions using RAPD markers. *Euphytica.* **115**: 7-16.
- Kidwell KK, Austinand DF and Osborn TC** (1994) RFLP evaluation of nine *Medicago* accessions representing the original germplasm source of North American alfalfa cultivars. *Crop Sci.* **34**: 230-236.
- Lai J, Yang W and Hisao J** (2001) An assessment of genetic relationship in cultivated tea clones and native wild tea in Taiwan using RAPD and ISSR markers. *Bot. Bull. Acad. Sin.* **42**: 93-100.

- Lakhanpaul S, Chanda S and Bhat KV** (2000) Random amplified polymorphic DNA (RAPD) analysis in Indian mungbean (*Vigna radiata* (L.) Wilczek) cultivars. *Genetica* **109**: 227-234.
- Mengoni A, Goriand A and Bazzicalupo M** (2000) Use of RAPD and microsatellite (SSR) variation to assess genetic relationships among population of tetraploid alfalfa, *Medicago sativa*. *Pl. Breed.* **119**: 311-317.
- Mishra RK, Chaudhury S, Ahmad A, Pradhan M and Siddiqi TO** (2009) Molecular analysis of tea clones (*Camellia sinensis*) using AFLP markers. *International Journal of Integrative Biology.* **5**(2): 130-136.
- Mishra RK and Mandi SS** (2004) Molecular profiling and development of DNA marker associated with drought tolerance in tea clones growing in Darjeeling. *Curr. Sci.* **87**(1): 60-66.
- Mondal TK, Singh HP and Ahuja PS** (2000) Isolation of genomic DNA from tea and other phenolic rich plants. *J. Planta Crops.* **28**(1): 30-34.
- Mondal TK, Bhattacharya A, Lakshmikumaran M and Ahuja PS** (2004) Recent advances of tea (*Camellia sinensis*) biotechnology. *Plant Cell Tissue and Organ Culture.* **76**: 195-254.
- Moonmoon S** (2006) Random amplified polymorphic DNA (RAPD) markers for genetic variation study among tomato varieties. MS Thesis. Dept. of Biotechnology. Bangladesh Agricultural University, Mymensingh. pp. 34-51.
- Mukhtar H and Ahmad N** (1999) Green tea in chemoprevention of cancer. *Toxicol. Sci.* **52**: 111-117.
- Mukhtar H and Ahmad N** (2000) Tea polyphenols-Prevention of cancer and optimizing health. *Amer. J. Clin. Nutr.* **71**: 1698S-1702S.
- Nei M** (1972) Genetic distance between populations. *American Nat.* **106**: 283-292.
- Orozco-Castillo C, KJ Chalmers, R Waugh and W Powell** (1994) Detection of genetic diversity and selective gene integration in coffee using RAPD markers. *TAG* **97**: 934-940.
- Paran I, Aftergoot E and Shifriess C** (1998) Variation in *Capsicum annum* revealed by RAPD and AFLP markers. *Euphytica* **9**(3): 167-173.
- Saini M, Singh S, Hussain Z and Yadav A** (2010) RAPD analysis in Mungbean (*Vigna radiata* (L.) Wilczek). Assessment of genetic diversity. *Indian J. Biotech.* **9**: 137-140.
- Sony SK, Habib MA and Islam MN** (2012) Genetic diversity analysis of thirteen mungbean (*Vigna radiata* (L.) Wilczek) cultivars using RAPD markers. *Bangladesh J. Bot.* **41**(2): 169-175.
- Sneath PHA and Sokal RR** (1973) Numerical taxonomy: Principles and Practice. Freeman, San Francisco, pp. 21-28.
- Tabassum N, Sony SK, bhajan SK and Islam MN** (2013) Analysis of genetic diversity in eleven tomato (*Lycopersicon esculentum* Mill.) varieties using RAPD markers. *Plant Tissue Cult. & Biotech.* **23**(1): 49-57.
- Virk PS, HJ Newbury, MT Jackson and BV Ford-Lloyd** (1995) The identification of duplicate accessions within Thirteen rice germplasm collection using RAPD analysis. *TAG* **90**: 1094-1055

**Wachira FN, Waugh R, Hackett CA and Powell W** (1995) Detection of genetic diversity in tea (*Camellia sinensis*) using RAPD markers. *Genome* **38**: 201-210.

**Weisburger JH** (1997) Tea and health: A historical perspective. *Cancer Lett.* **114**: 315-317.

**Welsh J and McClelland M** (1990) Fingerprinting genome using PCR with arbitrary primers. *Nucleic Acids Res.* **18**: 7213-7218.

**Williams JGK, Kubelik AR, Livak KJ Rafaliski JA and Tingey SV** (1990) DNA polymorphism amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Res.* **18**: 6531-6535.